

Neural Correlates of Human Echolocation of Path Direction during Walking

Katja Fiehler¹, Immo Schütz¹, Tina Meller¹, and Lore Thaler²

¹Department of Psychology, Justus-Liebig-University Giessen, Otto-Behaghel-Str. 10F,
35394 Giessen, Germany

²Department of Psychology, Durham University, South Road, Science Site, Lore Thaler,
Durham DH1 3LE, UK

Supplementary Tables

Table S1. Activations in the global RFX GLM for the sounds > baseline contrast. Results are cluster-level corrected at $z > 3.7$ with a cluster probability threshold of $p < 0.01$. Z-values reference peak activations within each cluster, corresponding peak coordinates are reported in MNI space (mm).

Voxels	p	z _{max}	X	Y	Z	L/R	Area(s)
298	3.00e-07	5.40	40	-26	12	R	Primary auditory cortex Planum temporale
297	3.11e-07	5.90	-44	-20	4	L	Primary auditory cortex Planum polare

Table S2. Activations in the global RFX GLM for the echo > no echo contrast. Results are cluster-level corrected at $z > 3.7$ with a cluster probability threshold of $p < 0.01$. Z-values reference peak activations within each cluster, corresponding peak coordinates are reported in MNI space (mm).

Voxels	p	z _{max}	X	Y	Z	L/R	Area(s)
501	1.89e-09	4.79	12	-76	50	R	Superior parietal lobule Anterior intra-parietal sulcus hIP2, hIP3 Inferior parietal lobule
180	7.99e-05	4.48	50	6	36	R	Premotor cortex Middle frontal gyrus
163	0.00016	4.64	66	-28	12	R	Primary auditory cortex Planum temporale Inferior parietal lobule
115	0.00133	4.42	-46	-26	6	L	Primary auditory cortex Planum temporale
84	0.00608	4.34	40	12	24	R	Inferior frontal gyrus