

Identification and quantification of polyphenols from *Cassia auriculata* L leaf, flower and flower bud using UPLC-QqQ-MS/MS

Devanesan Arul Ananth^{a,b}, Vijayaraghavan Mahalakshmi^c, Thilagar Sivasudha^d, Liron Klipcan^a and Zipora Tietel^a

^aGilat Research Center, Agricultural Research Organization, M.P. Negev 8531100, Israel; ^bCurrent affiliation: Department of Biotechnology, Karpagam Academy of Higher Education, Coimbatore - 641 021, Tamil Nadu, India; ^cDepartment of Biotechnology, Anna University, Chennai – 600 025, Tamil Nadu, India; ^dDepartment of Environmental Biotechnology, Bharathidasan University, Tiruchirappalli – 620 024, Tamil Nadu, India

CONTACT Zipora Tietel [**e-mail**] tietel@agri.gov.il

Supplementary material




Figure S1. UPLC-QqQ- MS/MS total ion chromatogram (TIC) of *C. auriculata* extracts.




Figure S2. Comparative quantitative analysis of polyphenolic classes contents across *C. auriculata* plant parts. Values are Mean \pm SE (n = 3).




Figure S3. Polyphenol contents of *C. auriculata*. (A) Across polyphenolic classes (B) Across plant parts. Values are Mean \pm SE (n = 3).