

Effects of the Belo Monte hydro-electric-dam complex on crocodilians in the Xingu River, Brazilian Amazonia

Zilca Campos¹, Fábio Muniz², William E. Magnusson^{3,**} and Guilherme Mourão^{1,***}

1 - Laboratório de vida selvagem, Embrapa Pantanal, Corumbá, Mato Grosso do Sul, MS 79320900, Brazil

2 - Laboratório de Evolução e Genética Animal (LEGAL), Universidade Federal do Amazonas (UFAM), Manaus, Amazonas, Brazil

3 - Instituto Nacional de Pesquisa da Amazônia (INPA), Manaus, Amazonas, Brazil

*Corresponding author; e-mail: zilca.campos@embrapa.br


**ORCID: 0000-0003-1988-3950

***ORCID: 0000-0002-8300-4191


Supplementary material

Figure S1. Distribution of the eight forest-sampling modules in relation to the river banks before reservoir filling (2014 and 2015) and after reservoir filling (2016 and 2017). Spatial distribution of individuals of *Paleosuchus trigonatus* (red triangles) and *Caiman crocodilus* (white circles) in the modules before (2014 and 2015) and after and after reservoir filling (2016 and 2017).


**Before flooding
2014-2015**



**After flooding
2016-2017**



**Before flooding
2014-2015**



**After flooding
2016-2017**

